arXiv:1201.3800v2 [math.CV] 1 Feb 2012

MAPPINGS OF FINITE DISTORTION OF POLYNOMIAL TYPE

CHANGYU GUO

Abstract. Suppose that $f:\mathbb{R}^n\to\mathbb{R}^n$ is a mapping of K-bounded p-mean distortion for some $p>n-1$. We prove the equivalence of the following properties of f: doubling condition for $J(x,f)$ over big balls centered at origin, boundedness of multiplicity function $N(f,\mathbb{R}^n)$, polynomial type of f and polynomial growth condition for f.

1. Introduction

Throughout this paper, we call a non-constant continuous mapping $f:\Omega\to\mathbb{R}^n$ a mapping of finite distortion if f belongs to the Sobolev space $W^{1,1}_{loc}(\Omega,\mathbb{R}^n)$ with $\frac{|Df|^n}{\log(e+|Df|)}\in L^1_{loc}(\mathbb{R}^n)$ and there exists a measurable function $K:\Omega\to[1,\infty)$ so that

$$|Df(x)|^n\le K(x)J(x,f), \tag{1}$$

for almost every $x\in\Omega$. Here $|Df(x)|$ and $J(x,f)$ are the operator norm and the Jacobian determinant of $Df(x)$, respectively. When $1\le K(x,f)\le K<\infty$ a.e., f is called a K-quasiregular mapping.

For a mapping of finite distortion f, the outer distortion function $K(\cdot,f)$ and the inner distortion function $K_I(\cdot,f)$ are defined as

$$K(x,f)=\frac{|Df(x)|^n}{J(x,f)}\quad and\quad K_I(x,f)=\frac{|D^{\#}f(x)|^n}{J(x,f)^{n-1}},$$

respectively, when $0<|Df(x)|,J(x,f)<\infty$, and $K(x,f)=K_I(x,f)=1$ otherwise. Here $|D^{\#}f(x)|$ is the adjoint matrix of $Df(x)$. Clearly

$$K_I(x,f)^{1/(n-1)}(x,f)\le K(x,f)\le K_I(x,f)^{n-1},$$

for almost every $x\in\Omega$.

We say that a mapping of finite distortion $f:\mathbb{R}^n\to\mathbb{R}^n$ has K-bounded p-mean distortion, $p\ge 1$, if there exists a constant $K\ge 1$ such that

$$\fint_{B(0,r)}K(x,f)^p dx\le K,\quad \forall 1\le r<\infty. \tag{2}$$

2000 *Mathematics Subject Classification.* 30C65.
Key words and phrases. mapping of finite distortion, modulus of path family.
C.Y.Guo was partially supported by the Academy of Finland grant 120972.

Let $f : \Omega \to \mathbb{R}^n$ be a mapping of finite distortion with $K(x, f) \in L^p_{loc}(\Omega)$ for some $p > n - 1$ and $E \subset \Omega$ a Borel set. We define the counting function

$$n(E, y) = \sum_{x \in f^{-1}(y) \cap E} i(x, f),$$

where $i(x, f)$ is the local index.

We let $A(a, r)$ be the average of $n(B(a, r), y)$ over $\overline{R}^n$ with respect to the n-dimensional spherical measure, i.e.

$$A(a, r) = \frac{2^n}{\omega_n} \int_{\mathbb{R}^n} \frac{n(B(a, r), y)}{(1 + |y|^2)^n} dy = \frac{2^n}{\omega_n} \int_{\overline{B}(a,r)} \frac{J_f(x)}{(1 + |f(x)|^2)^n} dx.$$

If a happens to be the origin, we simply denote it by $A(r) = A(0, r)$.

For a mapping of finite distortion $f : \mathbb{R}^n \to \mathbb{R}^n$, we define the lower order of f to be

$$\lambda_f = \liminf_{r \to \infty} \frac{\log A(r)}{\log r}. \tag{3}$$

We say that f has finite lower order if $\lambda_f < \infty$.

The main result of this paper is the following theorem that generalizes the corresponding statements for quasiregular mappings from [1].

Theorem A. *Let $f : \mathbb{R}^n \to \mathbb{R}^n$ be a mapping of K-bounded p-mean distortion for some $p > n - 1$. Then the following statements are equivalent*

- **1:** $J(x, f)$ *is doubling for balls centered at the origin with radius bigger than or equal to 1.*
- **2:** f *is of polynomial type, i.e.* $\lim_{x \to \infty} |f(x)| = \infty$ *and so* f *can be extended continuously to* $\overline{\mathbb{R}^n}$.
- **3:** $N(f, \mathbb{R}^n)$ *is bounded.*
- **4:** f *has polynomial growth, i.e. there exist positive constants* C_1 *and* C_2 *and integer* m *such that* $|f(x)| \leq C_1 |x|^m + C_2$.

Moreover, each of the above condition implies that $A(r)$ *is doubling for* $r \geq r_0$*, where* $r_0 > 0$ *is a constant. In particular,* f *has finite lower order.*

The lesson we draw from Theorem A is that ∞ must be a removable singularity if f does not grows too fast. Under the condition that f is of polynomial type, we actually conclude that f must cover the whole $\mathbb{R}^n$.

2. Preliminaries and Auxiliary results

2.1. **Area formula.** Let $f : \Omega \to \mathbb{R}^n$ be a mapping of finite distortion with distortion function $K(x, f) \in L^p_{loc}(\Omega)$ for some $p > n - 1$. Then f is sense-preserving, discrete and open by the result in [2], the latter meaning that $f^{-1}(y)$ cannot have accumulation points in Ω. In particular, $N(y, f, A) < \infty$ whenever $A \subset\subset \Omega$. Recall that $N(y, f, \Omega)$ is defined as the number of preimages of y under f in Ω. We also put $N(f,\Omega) := \sup_{y\in\mathbb{R}^n} N(y,f,\Omega)$.

An open, connected neighborhood $U \subset\subset \Omega$ of a point $x \in \Omega$ is called a normal neighborhood of x if $f(\partial U) = \partial f(U)$ and $U \cap f^{-1}(f(x)) = \{x\}$. Denote by $U(x,f,r)$ the x-component of $f^{-1}(B(f(x),r))$. For the following lemma, see Chapter I Lemma 4.9 in [8].

Lemma 1 (Lemma 4.9, [8]). *For each point* $x \in \Omega$ *there is* $\sigma_x > 0$ *such that* $U(x,f,r)$ *is a normal neighborhood of* x *whenever* $0 < r \leq \sigma_x$. *Moreover,* $\operatorname{diam} U(x,f,r) \to 0$ *as* $r \to 0$.

Next suppose that $x \in \Omega$ and $0 < r \leq \sigma_x$, where σ_x is as in the above lemma. The local index $i(x,f)$ of f at x is defined as

$$i(x,f) = N(f, U(x,f,r)).$$

Note in particular that $N(f,U(x,f,r))$ is independent of $r \leq \sigma_x$. Thus, $i(x,f) = 1$ if and only if $x \in \Omega\backslash B_f$, where B_f is the branch set of f. We refer the reader to [8] Chapter I for this discussion.

Suppose that $f : \Omega \to \mathbb{R}^n$ is a continuous, sense-preserving, discrete and open mapping. Let $\beta : [a,b) \to \mathbb{R}^n$ be a path and let $x \in f^{-1}(\beta(a))$. A path $\alpha : [a,c) \to \Omega$ is called a maximal f-lifting of β starting at x if $\alpha(a) = x$, $f \circ \alpha = \beta|_{[a,c)}$ and if whenever $c < c' \leq b$, there does not exist a path $\alpha' : [a,c) \to \Omega$ such that $\alpha = \alpha'|_{[a,c)}$ and $f \circ \alpha' = \beta|_{[a,c')}$.

Now let $x_1,\ldots,x_k$ be k different points of $f^{-1}(\beta(a))$ such that

$$m = \sum_{j=1}^{k} i(x_j,f).$$

We say that the sequence $\alpha_1,\ldots,\alpha_m$ of paths is a maximal sequence of f-liftings of β starting at the points $x_1,\ldots,x_k$ if each α_j is a maximal f-lifting of β such that

- $card\{j : \alpha_j(a) = x_i\} = i(x_i,f)$, $1 \leq i \leq k$,
- $card\{j : \alpha_j(t) = x\} \leq i(x,f)$ for all $x \in \Omega$ and all t.

The existence of maximal sequences of f-liftings is proved in Chapter II Theorem 3.2 in [8].

We close this subsection by the following well-known area formula.

Lemma 2. *Let* $f : \Omega \to \mathbb{R}^n$ *be a mapping of finite distortion with distortion function* $K(x,f) \in L^p_{loc}(\Omega)$ *for some* $p > n-1$ *and* η *be a non-negative Borel-measurable function on* $\mathbb{R}^n$. *Then*

$$\int_\Omega \eta(f(x))J(x,f)dx = \int_{\mathbb{R}^n} \eta(y)N(y,f,\Omega)dy. \tag{4}$$

The validity of the above lemma lies in the fact that f satisfies the so-called Lusin condition N. This is proven in [3] Theorem A. Recall that a mapping $f : \Omega \to \mathbb{R}^n$ is said to satisfy the Lusin condition N if the implication $|E| = 0 \Rightarrow |f(E)| = 0$ holds for all measurable sets $E \subset \Omega$.

2.2. **Basic modulus estimates.** We first give the definitions of the p-modulus and weighted p-modulus of a path family. Let E and F be subsets of $\overline{\Omega}$. We denote by $\Gamma(E,F,\Omega)$ the path family consisting of all locally rectifiable paths joining E to F in Ω. A Borel function $\rho\colon \mathbb{R}^n \to [0,\infty]$ is said to be admissible for $\Gamma(E,F,\Omega)$ if $\int_\gamma \rho\, ds \geq 1$ for all $\gamma \in \Gamma(E,F,\Omega)$. The p-modulus of a path family $\Gamma := \Gamma(E,F,\Omega)$ is defined as

$$\mathrm{mod}_p(\Gamma) := \inf\Big\{\int_\Omega \rho^p(x)\, dx : \rho\colon \mathbb{R}^n \to [0,\infty] \text{ is an admissible Borel function for } \Gamma\Big\}.$$

Let $\omega : \Omega \to [0,\infty]$ be a measurable function. The weighted p-modulus of the path family Γ is then defined as

$$\mathrm{mod}_{p,\omega}(\Gamma) := \inf\Big\{\int_\Omega \rho^p(x)\omega(x)\, dx : \rho\colon \mathbb{R}^n \to [0,\infty] \text{ is an admissible Borel function for } \Gamma\Big\}.$$

Note that when $\omega(x) = 1$ for all $x \in \Omega$, we recover the usual p-modulus mod_p. When $p = n$, we write mod_ω instead of $\mathrm{mod}_{n,\omega}$.

Next we introduce the p-modulus on spheres. Let Γ be a path family in $S(a,r)$, and let $1 < p < \infty$. Set

$$\mathrm{mod}_p^S(\Gamma) := \inf\Big\{\int_{S(a,r)} \rho^p(x)\, dS(x) : \rho\colon S(a,r) \to [0,\infty] \text{ is an admissible Borel function for } \Gamma\Big\}.$$

The notation Γ_f is used to denote the collection of all locally rectifiable paths in A having a closed subpath on which f is not absolutely continuous. The following K_o-inequality for mapping of finite distortion is due to Rajala [6].

Lemma 3 ([6], Theorem 2.1). *Let $f : \Omega \to \mathbb{R}^n$ be a mapping of finite distortion with $K(x,f) \in L^p_{loc}(\Omega)$ for some $p > n-1$. Let $A \subset \Omega$ be a Borel set with*

$$\sup_{y\in\mathbb{R}^n} N(y,f,A) < \infty,$$

and Γ a family of paths in A. If a function ρ is admissible for $f(\Gamma\backslash\Gamma_f)$, then

$$\mathrm{mod}_{K^{-1}(\cdot,f)}(\Gamma\backslash\Gamma_f) \leq \int_{\mathbb{R}^n} \rho^n(y) N(y,f,A)\,dy, \tag{5}$$

Moreover, $\mathrm{mod}_p(\Gamma_f) = 0$ *for all* $1 < p < n$.

The following important Väsälä inequality for mappings of finite distortion was proved by Koskela and Onninen [4]. Here $i(x,f)$ is the local index of f at a point x.

Lemma 4 ([4], Theorem 4.1)**.** *Let* $f : \Omega \to \mathbb{R}^n$ *be a mapping of finite distortion with* $K(x, f) \in L^p_{loc}(\Omega)$ *for some* $p > n-1$*. Let* Γ *be a path family in* Ω*,* Γ' *a path family in* $\mathbb{R}^n$*, and* m *a positive integer with the following property. For every path* $\beta : I \to \mathbb{R}^n$*, there are paths* $\alpha_1, \cdots, \alpha_m$ *in* Γ *such that* $f \circ \alpha_j \subset \beta$ *for all* j *and such that for every* $x \in \Omega$ *and* $t \in I$*,* $\alpha_j(t) = x$ *for at most* $i(x, f)$ *indices* j*. Then*

$$\operatorname{mod}(\Gamma') \leq \frac{\operatorname{mod}_{K_I(\cdot, f)}(\Gamma)}{m}. \tag{6}$$

For spherical rings, we prove the following upper bound for $K_I(\cdot, f)$-modulus.

Lemma 5. *Let* $f : \Omega \to \mathbb{R}^n$ *be a mapping of finite distortion with* $K(x, f) \in L^p_{loc}(\Omega)$ *for some* $p > n-1$*. Suppose that for some* $a \in \Omega$ *and* $R > 1$*,* $B(a, R) \subset\subset \Omega$ *and that*

$$I^s_a := \fint_{B(a,s)} K(x, f)^p dx \leq K < \infty, \quad \forall 1 \leq s \leq R.$$

There exists a constant $C > 0$*, depending only on* n *and* K*, such that*

$$\operatorname{mod}_{K_I(\cdot, f)}(\Gamma) \leq C \log^{1-n}(R/r). \tag{7}$$

for all $1 \leq 2r \leq R$*, where* Γ *is the family of all paths connecting* $\overline{B}(a, r)$ *and* $\mathbb{R}^n \backslash B(a, R)$*.*

Proof. Let $\rho(x) = \frac{1}{|x-a|} \log^{-1}(R/r)$ if $x \in B(a, r) \backslash \bar{B}(a, r)$ and $\rho(x) = 0$ otherwise. Then ρ is an admissible function for Γ. Let k be the smallest positive integer such that $2^k r \geq R$. Then it follows that

$$\begin{aligned}
\operatorname{mod}_{K_I(\cdot, f)}(\Gamma) &\leq \int_{B(a,R)\backslash \bar{B}(a,r)} \rho^n K_I(x, f) dx \\
&\leq \sum_{i=1}^{k} \int_{B(a,2^i r)\backslash \bar{B}(a,2^{i-1} r)} \frac{1}{|x-a|^n} \log^{-n}(R/r) K_I(x, f) dx \\
&\leq C(n) \log^{-n}(R/r) \sum_{i=1}^{k} K = C(n) K \log^{1-n}(R/r),
\end{aligned}$$

where, in the last inequality, we have used the condition $I^s_a \leq K$ for all $1 \leq s \leq R$. □

Next we will derive a lower bound estimate for the $K^{-1}(\cdot, f)$-modulus for spherical rings . The proof here is a small modification to the proof of Theorem 2.2 in [6].

Lemma 6. *Let* $f : \Omega \to \mathbb{R}^n$ *be a mapping of finite distortion with* $K(x, f) \in L^p_{loc}(\Omega)$ *some* $p > n-1$*. Let* E *and* F *be two continua. Assume that there exists a point* a *such that* $S(a, t)$ *intersects both* E *and* F *for* *all* $r' < t < R'$ *and* $B(a,R') \subset \Omega$. *Let* $Y = \fint_{B(a,R')\backslash B(a,r')} K(x,f)^p dx$. *Then we have the following lower bound:*

$$\mathrm{mod}_{K^{-1}(\cdot,f)}(\Gamma\backslash\Gamma_f) \geq C(n)\frac{R'-r'}{R'}Y^{-\frac{1}{p}}, \tag{8}$$

where Γ *is the family of paths joining* E *and* F *in* $B(a,R')\backslash\overline{B}(a,r')$.

Proof. Let ρ be an admissible function for $\mathrm{mod}_{K^{-1}(\cdot,f)}(\Gamma\backslash\Gamma_f)$ so that $\rho^n K^{-1}(\cdot,f)$ is integrable. Fix $p' = n - \frac{n}{p+1} \in (n-1,n)$. Then

$$\int_{S(a,t)} \rho^{p'}(x)dS(x) < \infty$$

for almost every $t \in (r',R')$. Now, by Lemma 3, $\mathrm{mod}_{p'}(\Gamma_f) = 0$ and thus $\mathrm{mod}^S_{p'}(\Gamma^t_f) = 0$ for almost all $t \in (r',R')$, where

$$\Gamma^t_f = \{\gamma \in \Gamma_f, |\gamma| \subset S(a,t)\}.$$

Lemma 2.4 in [6] gives for almost all $t \in (r',R')$,

$$\int_{S(a,t)} \rho^{p'}(x)dS(x) \geq \mathrm{mod}^S_{p'}(\Gamma^t\backslash\Gamma^t_f) = \mathrm{mod}^S_{p'}(\Gamma^t) \geq \frac{C_{p'}}{t^{p'+1-n}},$$

where $\Gamma^t = \{\gamma \in \Gamma : |\gamma| \subset S(a,t)\}$. We conclude that

$$1 \leq Ct^{p'+1-n}\int_{S(a,t)} \rho^{p'}(x)dS(x)$$

for almost every $t \in (r',R')$. Integrating over t, we get

$$(R'-r')essinf_{t\in(r',R')}\int_{S(a,t)} \rho^{p'}(x)dS(x) \leq \int_{B(a,R')\backslash B(a,r')} \rho^{p'}(x)dx.$$

Write $E = B(a,R')\backslash B(a,r')$. Then $|E|$ is bounded from above and below by dimensional constants times $(R'-r')R'^{n-1}$. Combining the above two inequalities gives

$$1 \leq CR'(|E|^{-1}\int_E \rho^{p'}(x)dx)^{1/p'}. \tag{9}$$

Writing $\rho^{p'}(x) = \rho^{p'}(x)K^{-p'/n}(x,f)K^{p'/n}(x,f)$ and using Holder's inequality, we see the righthand side of (9) is at most

$$CR'(|E|^{-1}\int_E \rho^n(x)K^{-1}(x,f)dx)^{1/n}(|E|^{-1}\int_E K^p(x,f)dx)^{1/np}. \tag{10}$$

Combining estimate (9) with (10), we obtain the desired estimate

$$C(n)\frac{R'-r'}{R'}Y^{-\frac{1}{p}} \leq \int_E \rho^n(x)K^{-1}(x,f)dx.$$

□

Lemma 7. *Let* $f : \mathbb{R}^n \to \mathbb{R}^n$ *be a mapping of* K*-bounded* p*-mean distortion for some* $p > n-1$. *Then* $f(\mathbb{R}^n)$ *cannot omit a set of positive* n*-capacity.*

Proof. Note that in our situation, all the requirements of Corollary 1.3 in [4] are satisfied and thus the conclusion follows. □

2.3. **Growth estimate.** In this subsection, we will derive a growth estimate on f under the assumption that $J(x, f)$ is doubling for balls centered at the origin with radius bigger than or equal to 1. The idea of the proof is from Lemma 12.3 in [5]. For the statement of the following results, we introduce the notations $L(0,R) = \sup\{|f(x)-f(y)| : |x-y| \le R\}$ and $l(0,R) = \inf\{|f(x) - f(y)| : |x-y| \ge R\}$.

Lemma 8. *Let* $f : \mathbb{R}^n \to \mathbb{R}^n$ *be a non-constant mapping of* K*-bounded* p*-mean distortion for some* $p > n-1$. *If* $J(x, f)$ *is* C*-doubling for all balls centered at the origin with radius bigger than or equal to 1, then there exist* $C' = C'(n,K) > 0$ *and* $R_0 > 0$ *such that*

$$L(0,R) \le C'R^k \tag{11}$$

for all $R > R_0$, *where* $k = \frac{\log_2 C+1}{n}$.

Proof. By Lemma 8, we may set R_0 to be such a number that $L(0,R) = 1$. Let $R > R_0$. We fix a point $a \in \overline{B}(0,R)$ such that

$$|f(a) - f(0)| = L(0,R). \tag{12}$$

By Lemma 7, we may choose $\gamma : [0,\infty) \to \mathbb{R}^n \backslash B(f(0), L(0,R))$, starting at $f(a)$ with $|\gamma|$ unbounded and a maximal f-lifting of γ' of γ starting at a. Then $|\gamma'|$ is also unbounded. Lemma 6 implies that

$$\mathrm{mod}_{K^{-1}(\cdot,f)}(\Gamma\backslash\Gamma_f) \ge C(n,K)R^{-1}, \tag{13}$$

where Γ is the family of curves joining $\overline{B}(0,1)$ to $|\gamma'| \cap \overline{B}(2R)$ in $B(0,2R)$. Since every path $\eta \in f(\Gamma)$ intersects $f(\overline{B}(0,1))$ and $\mathbb{R}^n \backslash B(f(0), L(0,R))$, the function $\rho : \mathbb{R}^n \to [0,\infty]$ defined by

$$\rho(y) = \frac{2}{L(0,R)} \chi_{f(B(0,2R))}(y)$$

is a test function for $f(\Gamma)$. By Lemma 2,

$$\int_{\mathbb{R}^n} N(y, f, B(0,2R))\rho^n(y)dy = \frac{2^n}{L(0,R)^n} \int_{B(0,2R)} J(x,f)dx \le C_1 \frac{R^s}{L(0,R)^n} \tag{14}$$

for $s = \log_2 C$. The claim now follows from Lemma 3, (13) and (14). □

Corollary Let $f : \mathbb{R}^n \to \mathbb{R}^n$ be a non-constant mapping of K-bounded p-mean distortion for some $p > n-1$. If $A(r)$ is C-doubling for $r \ge 1$ and $L(0,r) \le C_1 l(0,r)$ for $r > r_0$, then there exist $C' = C'(n,K) > 0$ and $R_0' > 0$ such that

$$l(0,2R) \ge C'R^{\alpha} \tag{15}$$

for all $R > R_0$, where $\alpha = \frac{1}{n(\log_2 C+1)}$. In particular, $f(\mathbb{R}^n)$ contains balls of arbitrary large radius.

Proof. The proof is quite similar to Lemma 8. Let R_0 be the same number as in the proof of Lemma 8 and $R'_0 = \max\{r_0, R_0\}$. For $R > R'_0$ we write

$$\rho(y) = \frac{2}{l(0,2R)}\chi_{f(B(0,2R))}(y)$$

to be the test function for $f(\Gamma)$ and

$$\begin{aligned}\int_{\mathbb{R}^n} N(y,f,B(0,2R))\rho^n(y)dy &= \frac{2^n}{l(0,2R)^n}\int_{B(0,2R)} J(x,f)dx \\ &\leq \frac{2^n(1+L(0,2R)^2)^n}{l(0,2R)^n}\int_{B(0,2R)}\frac{J(x,f)}{(1+|f(x)|^2)^n}dx \\ &\leq C(n)l(0,2R)^nR^s\end{aligned}$$

for $s = \log_2 C$. □

Remark. This result can be viewed as a covering theorem for mappings of finite distortion, comparing it with Corollary 1.2 in [7].

3. Proof of theorem A

Proof. The implication "**3 ⇒ 1**" follows from modulus estimates. First of all, Lemma 2 implies that

$$\int_{2B} J(x,f)dx = \int_{f(2B)} N(y,f,2B)dy \leq N(f,\mathbb{R}^n)|f(2B)|$$

and

$$\int_{B} J(x,f)dx = \int_{f(B)} N(y,f,B)dy \geq |f(B)|.$$

Therefore it suffices to show that

$$|f(2B)| \leq C|f(B)|$$

whenever $B \subset \mathbb{R}^n$ is a ball centered at origin with radius bigger than or equal to 1.

To this end, fix a ball $B = B(0,\eta) \subset \mathbb{R}^n$ with $\eta \geq 1$. Since f is open, $f(0)$ is an interior point of the open connected set $U = f(B)$. Let $d = \mathrm{dist}(f(0), \partial U)$ and set

$$r = \max_{y\in\partial U'}|f(0) - y|,$$

where $U' = f(2B)$. Then clearly $2r \geq \mathrm{diam}(U')$. Let L_0 be the line segment of length d joining $f(0)$ to ∂U and let L_1 be a half line in $\mathbb{R}^n$ joining a point in $\partial B(f(0),r) \cap \partial U'$ to ∞ in $\mathbb{R}^n \cup \{\infty\}\backslash B(f(0),r)$. Then some lifts of L_0 and L_1 join 0 to ∂B, and $\partial 2B$ to ∞, respectively, and hence by Lemma 6

$$\mathrm{mod}_{K^{-1}(\cdot,f)}(\Gamma\backslash\Gamma_f) \geq C > 0,$$

where Γ is the family of paths joining $f^{-1}(L_0)$ and $f^{-1}(L_1)$ in $3B$. On the other hand, Lemma 3 gives us

$$\operatorname{mod}_{K^{-1}(\cdot,f)}(\Gamma\backslash\Gamma_f) \leq N(y,f,\mathbb{R}^n)\operatorname{mod}(L_0,L_1,\mathbb{R}^n) \leq C(\log\frac{r}{d})^{1-n}.$$

We thus obtain $r \leq Cd$, which implies that $f(B)$ and $f(2B)$ have comparable volumes, as desired.

Next, we prove the equivalence of **2** and **3**. First we show that **2** implies **3**. To this end, fix a point $y \in f(\mathbb{R}^n)$. Since f is discrete and $\lim_{x\in\infty}|f(x)| = \infty$, there exists a ball $B = B(0,r) \subset \mathbb{R}^n$ such that

$$N(y,f,\mathbb{R}^n) = N(y,f,B) \leq \sum_{x\in f^{-1}(y)} i(x,f) =: M < \infty. \tag{16}$$

Now suppose that there exists a point $v \in X$ with

$$\sum_{x\in f^{-1}(v)} i(x,f) > M,$$

and choose a compact path γ beginning at v and ending at y. Then there are at least $M+1$ lifts γ^j of γ starting at $f^{-1}(v)$, and each of them either ends at some $x \in f^{-1}(y)$ or leaves every compact subset of $\mathbb{R}^n$. The latter cannot happen for any j since f is of polynomial type. But, by (16), the former can occur for at most M of the lifts γ^j. This is a contradiction and thus **2** implies **3**.

Now assume **3** and suppose that **2** does not hold. Then there exists a sequence (a_i) of points in $\mathbb{R}^n$, such that $|a_i|$ increases to infinity but

$$\limsup_{i\to\infty}|f(a_i) - f(0)| = R < \infty. \tag{17}$$

Passing to a subsequence if necessary, we may assume that

$$\operatorname{diam} f(B(0,|a_i|)) \geq R/2$$

for every $i \in \mathbb{N}$. Note that f is continuous and open, and hence it is monotone. It follows that

$$\operatorname{diam} f(S(0,|a_i|)) \geq R/2. \tag{18}$$

Since $N(f,\mathbb{R}^n) < \infty$, we may fix a $\delta > 0$ so that $U(x_j,\delta)$ is a normal neighborhood of x_j for every $x_j \in f^{-1}(f(0))$. Then

$$\cup_{x_j\in f^{-1}(f(0))} U(x_j,\delta) \subset B(0,t) \tag{19}$$

for some $t > 0$. When $|a_i| > t$, let Γ_i be the family of all paths joining $B(f(0),\delta)$ and $f(S(0,|a_i|))$ in $\mathbb{R}^n$. Then by (17), (18) and the n-Loewner property of $\mathbb{R}^n$, there exists a constant $C > 0$ such that $\operatorname{mod}(\Gamma_i) \geq C$ for every i. Denote by Γ_i' the family of all lifts γ' of $\gamma \in \Gamma_i$ starting at $S(0,|a_i|)$.

By (19), every $\gamma' \in \Gamma_i'$ either intersects $B(0,t)$ or leaves every compact set in $\mathbb{R}^n$. The latter family of paths have $K_I(\cdot,f)$-modulus zero. All other paths start at $S(0,|a_i|)$ and intersect $S(0,t)$, so

$$\operatorname{mod}_{K_I(\cdot,f)}(\Gamma_i') \leq C(n,K,t)\log^{1-n}(|a_i|/t) \to 0, \quad as\ i \to \infty$$

by the remark after Lemma 5. We thus get a contradiction to Lemma 4. Therefore, **2** follows from **3**.

Now we show the implication "**1** $\Rightarrow$ **3**". First we suppose R is large and fix a point $y \in f(B(0,R))$ with m preimage points $x_1, \cdots, x_m$ inside $B(0,R) \subset \mathbb{R}^n$, where m is a large positive integer to be estimated later. We choose $\delta > 0$ so that $U(x_i, \delta) \subset B(0,R)$ is a normal neighborhood for each $i = 1, \cdots, m$, and the sets $U(x_i, \delta)$ are pairwise disjoint.

By Lemma 7, we can choose a point $f(q) \in f(\mathbb{R}^n)$ with $|y - f(q)|$ as large as desired, and a path

$$\gamma : [0, \infty) \to \mathbb{R}^n \backslash B(y, |y - f(q)|),$$

starting at $f(q)$, such that $|\gamma|$ is unbounded. Then

$$\operatorname{mod}(\Gamma) = \operatorname{mod}(\overline{B}(y,\delta), |\gamma|, \mathbb{R}^n) \geq C_1 (\log \frac{|y - f(q)|}{\delta})^{1-n}, \tag{20}$$

where $C_1 > 0$ does not depend on q.

By Lemma 8 and our choice of γ, there exist $\alpha > 0$ such that

$$\operatorname{dist}(0, f^{-1}(|\gamma|)) \geq C|y - f(q)|^{\alpha} \tag{21}$$

when the right-hand term is larger than some fixed constant. For each $\eta \in \Gamma$, there are at least m maximal f-liftings η_i starting at the points $x_i \in B(0,R)$. Moreover, by the above estimate, each of them intersects $\mathbb{R}^n \backslash B(0, C|y - f(q)|^{\alpha})$. Denote the family of all such lifts by Γ'. Then by Lemma 4 and Lemma 5,

$$\operatorname{mod}(\Gamma) \leq \frac{\operatorname{mod}_{K_I(\cdot,f)}(\Gamma')}{m} \leq \frac{C_1}{m} \log^{1-n}(C_2|y - f(q)|^{\alpha}/R). \tag{22}$$

Combining (20) and (22) yields

$$|y - f(q)|^{\alpha} \leq \frac{R}{\delta} |y - f(q)|^{\frac{C}{m^{1/(n-1)}}}. \tag{23}$$

We conclude from (23) that $m \leq (\frac{C}{\alpha})^{n-1}$. To see this, suppose that the conclusion is not true, i.e. $m > (\frac{C}{\alpha})^{n-1}$. We may let $|y - f(q)|$ be as large as we wish and obtain a contradiction with (23).

The implication "**1** $\Rightarrow$ **4**" follows immediately from Lemma 8. The proof of implication "**4** $\Rightarrow$ **3**" is the same as that in the implication "**1** $\Rightarrow$ **3**". Indeed, (21) follows immediately from Lemma 8.

Finally we only need to show that one of the above four conditions implies that $A(r)$ is doubling for $r \geq r_0$ for some positive constant r_0. We want to prove that **3** implies that $A(r)$ is doubling for all $r \geq r_0$. The proof is similar to the proof for the implication "**3** $\Rightarrow$ **1**". Let $B = B(0,r)$ with $r \geq r_0$, where $r_0 > 0$ is to be determined later. Lemma 2 gives us

$$\begin{aligned} A(2r) &= \int_{2B} \frac{J(x,f)}{(1 + |f(x)|^2)^n} dx \\ &= \int_{f(2B)} \frac{N(y,f,2B)}{(1 + |y|^2)^n} dy \leq N(f, \mathbb{R}^n) \int_{f(2B)} \frac{1}{(1 + |y|^2)^n} dy \end{aligned}$$

and

$$A(r)=\int_B \frac{J(x,f)}{(1+|f(x)|^2)^n}dx=\int_{f(B)}\frac{N(y,f,2B)}{(1+|y|^2)^n}dy\geq\int_{f(B)}\frac{1}{(1+|y|^2)^n}dy.$$

Therefore it suffices to show that

$$\int_{f(2B)}\frac{1}{(1+|y|^2)^n}dy\leq C\int_{f(B)}\frac{1}{(1+|y|^2)^n}dy.$$

Then we follow the proof of implication of "**3** $\Rightarrow$ **1**". Since f is open, $f(0)$ is an interior point of the open connected set $U=f(B)$. Let $d=\mathrm{dist}(f(0),\partial U)$ and set

$$r'=\max_{y\in\partial U'}|f(0)-y|,$$

where $U'=f(2B)$. Now we choose r_0 so that $r\geq r_0$ implies that $d\geq 2$. Arguing as before, we conclude that $r'\leq Cd$. Thus we have

$$\int_{f(B)}\frac{1}{(1+|y|^2)^n}dy\geq\int_{B(f(0),d)}\frac{1}{(1+|y|^2)^n}dy\geq C(n)\int_0^d\frac{t^{n-1}}{(1+t^2)^n}dt.$$

On the other hand,

$$\int_{f(2B)}\frac{1}{(1+|y|^2)^n}dy\leq\int_{B(f(0),r')}\frac{1}{(1+|y|^2)^n}dy\leq C(n)\int_0^{r'}\frac{t^{n-1}}{(1+t^2)^n}dt.$$

Note that if we split the integral $\int_0^l\frac{t^{n-1}}{(1+t^2)^n}dt$ into two parts,

$$\int_0^l\frac{t^{n-1}}{(1+t^2)^n}dt=\int_0^1\frac{t^{n-1}}{(1+t^2)^n}dt+\int_1^l\frac{t^{n-1}}{(1+t^2)^n}dt,$$

then, for the second integral, we have that

$$\frac{1}{n\cdot 2^n}(1-\frac{1}{l^n})\leq\int_1^l\frac{t^{n-1}}{(1+t^2)^n}dt\leq\frac{1}{n}(1-\frac{1}{l^n}).$$

Since $1-\frac{1}{d^n}\geq 1-\frac{1}{2^n}$ and $1-\frac{1}{r^n}\leq 1$, we conclude that $A(r)$ is doubling for $r\geq r_0$. ☐

Remark. Note that the boundedness of $N(f,\mathbb{R}^n)$ does not imply that $J(x,f)$ being doubling for all balls centered at the origin as one may consider the mapping

$$F(x)=\begin{cases}\frac{x}{|x|}\log^{-1}\frac{1}{|x|} & \text{if } |x|\leq 1/e\\ x & \text{if } |x|\geq 1/e.\end{cases}$$

Acknowledgements

We thank Academic Professor Pekka Koskela for many useful suggestions and for carefully reading the manuscript. We also thank Academy Research Fellow Kai Rajala for many valuable discussions.

REFERENCES

[1] J.Heinonen and P.Koskela, Weighted Sobolev and Poincare inequalities and quasiregular mappings of polynomial type, Math. Scand. 77 (1995), no. 2, 251-271.

[2] J.Kauhanen, P.Koskela and J.Maly, Mappings of finite distortion: discreteness and openness, Arch. Ration. Mech. Anal. 160 (2001), no. 2, 135-151.

[3] J.Kauhanen, P.Koskela and J.Maly, Mappings of finite distortion: condition N, Michigan Math. J. 49 (2001), no. 1, 169-181.

[4] P.Koskela and J.Onninen, Mappings of finite distortion: capacity and modulus inequalities, J. Reine Angew. Math. 599 (2006), 1-26.

[5] J.Onninen and K.Rajala, Quasiregular mappings to generalized manifolds, J. Anal. Math. 109 (2009), 33-79.

[6] K.Rajala, Mappings of finite distortion: the Rickman-Picard theorem for mappings of finite lower order, J. Anal. Math. 94 (2004), 235-248.

[7] K.Rajala, Bloch's theorem for mappings of bounded and finite distortion, Math. Ann. 339 (2007), no. 2, 445-460.

[8] S.Rickman, Quasiregular mappings. Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], 26. Springer-Verlag, Berlin, 1993, x+213 pp.

(Changyu Guo) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ, P.O. BOX 35, FI-40014 UNIVERSITY OF JYVÄSKYLÄ, FINLAND
E-mail address: `changyu.c.guo@jyu.fi`